

Competing sexual and asexual generic names in *Pucciniomycotina* and *Ustilaginomycotina* (*Basidiomycota*) and recommendations for use

M. Catherine Aime¹, Lisa A. Castlebury², Mehrdad Abbasi¹, Dominik Begerow³, Reinhard Berndt⁴, Roland Kirschner⁵, Ludmila Marvanová⁶, Yoshitaka Ono⁷, Mahajabeen Padamsee⁸, Markus Scholler⁹, Marco Thines¹⁰, and Amy Y. Rossman¹¹

¹Purdue University, Department of Botany and Plant Pathology, West Lafayette, IN 47901, USA; corresponding author e-mail: maime@purdue.edu

²Mycology & Nematology Genetic Diversity and Biology Laboratory, USDA-ARS, Beltsville, MD 20705, USA

³Ruhr-Universität Bochum, Geobotanik, ND 03/174, D-44801 Bochum, Germany

⁴ETH Zürich, Plant Ecological Genetics, Universitätstrasse 16, 8092 Zürich, Switzerland

⁵Department of Biomedical Sciences and Engineering, National Central University, 320 Taoyuan City, Taiwan

⁶Czech Collection of Microorganisms, Faculty of Science, Masaryk University, 625 00 Brno, Czech Republic

⁷Faculty of Education, Ibaraki University, Mito, Ibaraki 310-8512, Japan

⁸Systematics Team, Manaaki Whenua Landcare Research, Auckland 1072, New Zealand

⁹Staatliches Museum f. Naturkunde Karlsruhe, Erbprinzenstr. 13, D-76133 Karlsruhe, Germany

¹⁰Senckenberg Gesellschaft für Naturforschung, Frankfurt (Main), Germany

¹¹Department of Botany & Plant Pathology, Oregon State University, Corvallis, OR 97333, USA

Abstract: With the change to one scientific name for pleomorphic fungi, generic names typified by sexual and asexual morphs have been evaluated to recommend which name to use when two names represent the same genus and thus compete for use. In this paper, generic names in *Pucciniomycotina* and *Ustilaginomycotina* are evaluated based on their type species to determine which names are synonyms. Twenty-one sets of sexually and asexually typified names in *Pucciniomycotina* and eight sets in *Ustilaginomycotina* were determined to be congeneric and compete for use. Recommendations are made as to which generic name to use. In most cases the principle of priority is followed. However, eight generic names in the *Pucciniomycotina*, and none in *Ustilaginomycotina*, are recommended for protection: *Classicula* over *Naiadella*, *Gymnosporangium* over *Roestelia*, *Helicobasidium* over *Thanatophyllum* and *Tuberculina*, *Melampsorella* over *Peridermium*, *Milesina* over *Milesia*, *Phragmidium* over *Aregma*, *Sporobolomyces* over *Blastoderma* and *Rhodomyces*, and *Uromyces* over *Uredo*. In addition, eight new combinations are made: *Blastospora juruensis*, *B. subneurophylla*, *Cronartium bethelii*, *C. kurilense*, *C. sahoanum*, *C. yamabense*, *Milesina polypodii*, and *Prospodium crusculum* combs. nov.

Key words:

Basidiomycetes
pleomorphic fungi
taxonomy
protected names
rejected names
unit nomenclature
new taxa

Article info: Submitted: 16 March 2018; Accepted: 26 March 2018; Published: 26 April 2018.

INTRODUCTION

With the change to one scientific name for all *Fungi* in accordance with the *International Code of Nomenclature for algae, fungi, and plants* (ICN, McNeill *et al.* 2012), it has been necessary to determine which generic name should be applied when two or more generic names are typified by different morphs of the same genus. Recommendations about which generic name to use have been made primarily by specialized Subcommissions or Working Groups of the International Commission for the Taxonomy of Fungi (ICTF), each representing a different group of *Fungi*. A number of papers recommending generic names for use have been published including the major groups of *Sordariomycetes* such as *Diaporthales* (Rossman *et al.* 2015a), *Hypocreales* (Rossman *et al.* 2013, Quandt *et al.* 2014, Kepler *et al.* 2017), *Magnaporthales* (Zhang *et al.* 2016), *Microascales* and

Ophiostomatales (de Beer *et al.* 2013), *Xylariaceae* (Stadler *et al.* 2013) and remaining *Sordariomycetes* (Réblová *et al.* 2016) as well as *Dothideomycetes* (Rossman *et al.* 2015b), *Eurotiales* (Samson *et al.* 2014, Visagie *et al.* 2014), *Leotiomycetes* including *Erysiphales* (Braun 2013, Johnston *et al.* 2014), *Orbiliomycetes* (Baral *et al.* 2017), *Pezizomycetes* (Healy *et al.* 2016), yeasts (Daniel *et al.* 2014) and overlooked generic names in the *Ascomycota* (Rossman *et al.* 2016). A companion paper to this one on the *Agaricomycotina* is in progress. Generic names that follow the principle of priority whether they are sexually or asexually typified do not need to be confirmed by the Nomenclature Committee for Fungi (NCF) and General Committee of the International Botanical Congress (IBC). The former Art. 57.2 which required approve for the use of asexually typified names over sexually typified ones was removed from the *Code* at the 2017 International Botanical Congress (Turland *et al.* 2017, 2018). However,

© 2018 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

Attribution: You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

Non-commercial: You may not use this work for commercial purposes.

No derivative works: You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

those generic names that do not have priority must be approved by the NCF.

Generic names of *Pucciniomycotina* and *Ustilaginomycotina* (*Basidiomycota*), representing sexually and asexually typified genera that compete for use have been evaluated by the *Heterobasidiomycetes* Working Group and the International Subcommittee on Rust Taxonomy of the ICTF. Members of these groups have been consulted and agree with the recommendations made herein about which generic name to use¹. The comprehensive list of sexual-aseexual generic names by Wijayawardene *et al.* (2012) was used as the initial starting point for discovering competing basidiomycete generic names. Citations for generic names and type species are based on *Index Fungorum*. Each set of generic names was evaluated based on the phylogenetic placement of their type species using the literature to determine whether the names are congeneric. Then a recommendation for use is made considering such factors as priority, number of species and thus number of required name changes, use outside of the ICN, frequency of citations in the literature and frequency of use based primarily on number of citations in Google Scholar (GS).

Here we advocate abandoning the use of many asexually typified generic names in *Pucciniales* such as *Aecidium*, *Caeoma*, *Milesia*, *Peridermium*, *Roestelia*, and *Uredo* for their corresponding sexually typified generic names. This is problematic as many species have been described using these names. Often it is difficult to determine in which genus a rust fungus belongs because of the lack of the telial stage. However, these formal asexually typified generic names are based on type species that are synonyms of species in other genera. In the case of *Aecidium* and *Caeoma* these generic names are synonyms of an earlier generic name, *Puccinia*. For *Peridermium* the type species is a synonym of a species in *Melamporella*, a later generic name that is recommended for protection over *Peridermium*; most species described in *Peridermium*, however, belong in *Cronartium*. Similarly, the type species of the generic name *Roestelia* is a synonym of the type species in *Gymnosporangium*, an important and well-defined genus that is recommended for protection. Finally, the generic name *Uredo* is a synonym of the later name *Uromyces*, which is recommended for use for two reasons. Firstly, species of *Uredo* belong to many different rust genera, and secondly *Uromyces* is a commonly used generic name. See under those generic names for more detailed discussion.

Although the abandoned *Pucciniales* asexual morph names should no longer be used to describe new species, the species already described in those names are not invalidated and can still be used. As for all scientific names

that cannot be correctly placed in a genus, the problem with the generic name could be noted by including quotes around the name or by indicating the temporary taxonomic placement with "pro tem." as has been suggested for anamorphically typified species of *Ustilaginomycotina* (Wang *et al.* 2015b). Efforts will be made in the future to determine the correct generic placement for the names currently in use in these asexually typified genera. With the increasingly available use of molecular data for determining phylogenetic relationships, the need for these asexually typified names of rust fungi should be eliminated or, as has been done for years, species known only from an asexual state can be placed in the genus most likely to be considered correct.

Generic names used for asexual rust fungi can, however, be used in a non-latinized sense to describe a morphology, as illustrated in Cummins & Hiratsuka (2003). As in other groups of fungi, for example, a species may be described as having ascomata that are nectria-like, meaning they produce light- to bright-coloured, fleshy perithecia, or fusarium-like meaning that they produce fusiform, multiseptate, hyaline conidia. Thus, one can refer to an asexual morph of a rust fungus as being aecidium-like, caeoma-like, peridermium-like, or uredo-like.

Twenty-one sets of sexual and asexual generic names in *Pucciniomycotina* and eight sets of generic names in *Ustilaginomycotina* were determined to compete for use and evaluated for recommended use. Most of the generic names recommended for use have priority. None of the names of *Ustilaginomycotina* but eight generic names of *Pucciniomycotina* do not have priority and are recommended for protection over older names, specifically: *Classicula* over *Naiadella*, *Gymnosporangium* over *Roestelia*, *Helicobasidium* over *Thanatophytum* and *Tuberculina*, *Melampsorella* over *Peridermium*, *Milesina* over *Milesia*, *Phragmidium* over *Aregma*, *Sporobolomyces* over *Rhodomyces* and *Blastoderma*, and *Uromyces* over *Uredo*. Details about each set of generic names and the basis for each decision are presented below with an S indicating a sexually typified name and an A indicating an asexually typified name. The citation for each generic name, type species, accepted name of type species, and action required are listed in Table 1 for *Pucciniomycotina* and Table 2 for *Ustilaginomycotina*.

COMPETING SEXUAL AND ASEQUAL GENERIC NAMES OF *Pucciniomycotina*, *Basidiomycota* AND RECOMMENDATIONS FOR USE

Use *Basidiopycnis* Oberw. *et al.* 2006 (S) rather than *Basidiopycnides* J. Reid *et al.* 2008 (A)

The monotypic generic name *Basidiopycnis* is typified by the sexual species *B. hyalina*, described from bark beetle galleries (Oberwinkler *et al.* 2006). Another monotypic generic name *Basidiopycnides* typified by *B. albertensis* was described for an asexual morph also from bark beetle galleries (Hausner *et al.* 2008). Kirschner & Oberwinkler (2009) confirmed the relationship between these types as sexual and asexual morphs of the same species and

¹*Heterobasidiomycetes* Working Group: Dominik Bergerow, Convenor; Cathie Aime, Stephan Helfer, David McLaughlin, Hai Nguyen, Maj Padamsee, Angela Schäfer, Qi-Ming Wang, and Andrey Yurkov.

International Subcommittee on Rust Taxonomy: Cathie Aime, Chair; Mehrdad Abbasi, Gaddam Bagyanarayana, Reinhard Berndt, Cvetomir Denchev, José Dianese, Richard Hamelin, Stephan Helfer, Drew Minnis, Yoshitaka Ono, and Markus Scholler.

Table 1. Recommended generic names of *Pucciniomycotina* among sexual-aseexual names that compete for use. The recommended name is in bold; see text for the rationale behind these recommendations. For each generic name this list provides the author, its date and place of publication, type species of the genus, its basionym, their dates of publication and the currently accepted name, if different. The action required is indicated in the last column such as approval by the Nomenclature Committee for Fungi (NCF) for those generic names that do not have priority and thus need protection.

Recommended generic name	Synonymous alternate morph generic name(s)	Action required
Basidiopycnis Oberw. <i>et al.</i> in <i>Mycologia</i> 98 : 639. 2006. Type: <i>B. hyalina</i> Oberw. <i>et al.</i> 2006.	<i>Basidiopycnides</i> J. Reid <i>et al.</i> in <i>Mycotaxon</i> 103 : 285. 2008. Type: <i>B. albertensis</i> J. Reid <i>et al.</i> 2008, now regarded as <i>Basidiopycnis hyalina</i> Oberw. <i>et al.</i> 2006.	None.
Blastospora Dietel in <i>Ann. Mycol.</i> 6 : 222. 1908. Type: <i>B. smilacis</i> Dietel 1908.	<i>Pelastoma</i> M. Salazar <i>et al.</i> in <i>Mycologia</i> 104 : 116. 2012. Type: <i>P. juruensis</i> (Henn.) M. Salazar <i>et al.</i> 2012, basionym: <i>Uredo juruensis</i> Henn. 1904, now regarded as <i>Blastospora juruensis</i> (Henn.) Aime & Rossman 2018.	None.
Camptobasidium Marvanová & Suberkr. in <i>Mycologia</i> 82 : 209. 1990. Type: <i>C. hydrophilum</i> Marvanová & Suberkr. 1990.	<i>Crucella</i> Marvanová & Suberkr., <i>Mycologia</i> 82 : 212, 1990. Type: <i>C. subtilis</i> Marvanová & Suberkr. 1990, now regarded as <i>Camptobasidium hydrophilum</i> Marvanová & Suberkr. 1990.	None.
Classicula R. Bauer <i>et al.</i> in <i>Mycologia</i> 95 : 757. 2003. Type: <i>C. fluitans</i> <i>et al.</i> 2003.	<i>Naiadella</i> Marvanová & Bandoni in <i>Mycologia</i> 79 : 579. 1987. Type: <i>N. fluitans</i> Marvanová & Bandoni 1987, now regarded as <i>Classicula fluitans</i> R. Bauer <i>et al.</i> 2003.	Protection needed by NCF.
Cronartium Fr., <i>Observ. Mycol.</i> 1 : 220. 1815. Type: <i>C. asclepiadeum</i> (Willd.) Fr. 1815, basionym: <i>Erineum asclepiadeum</i> Willd. 1806, now regarded as <i>Cronartium flaccidum</i> (Alb. & Schwein.) G. Winter 1880.	<i>Endocronartium</i> Y. Hirats. in <i>Canad. J. Bot.</i> 47 : 1493. 1969. Type: <i>E. harknessii</i> (J.P. Moore) Y. Hirats. 1969, basionym: <i>Peridermium harknessii</i> J.P. Moore 1884, now regarded as <i>Cronartium harknessii</i> (J.P. Moore) E. Meinecke 1920.	None.
Elateraecium Thirum. <i>et al.</i> in <i>Mycologia</i> 58 : 394. 1966. Type: <i>E. salaciicola</i> Thirum. <i>et al.</i> 1966.	<i>Hiratsukamyces</i> Thirum. <i>et al.</i> <i>Sydowia</i> 27 : 80, 1975. Type: <i>H. salaciicola</i> Thirum. <i>et al.</i> 1975, now regarded as <i>Elateraecium salaciicola</i> Thirum. <i>et al.</i> 1966.	None.
Gerwasia Racib. in <i>Bull. Int. Acad. Sci. Cracovie, Cl. Sci. Math.</i> 3 : 270. 1909. Type: <i>G. rubi</i> Racib. 1909.	<i>Morispota</i> Salazar-Yepes <i>et al.</i> in <i>Caldasia</i> 29 : 108. 2007. Type: <i>M. tenella</i> (H.S. Jacks. & Holw.) Salazar-Yepes <i>et al.</i> 2007, basionym: <i>Mainsia tenella</i> H.S. Jacks. & Holw. 1931, now regarded as <i>Gerwasia tenella</i> (H.S. Jacks. & Holw.) Buriticá 1994.	None.
Gymnosporangium R. Hedw. ex DC., in Lamarck & de Candolle, <i>Fl. franç.</i> , ed. 3 2: 216. 1805. Type: <i>G. fuscum</i> DC. 1805, now regarded as <i>Gymnosporangium sabinae</i> (Dicks.) G. Winter 1884.	<i>Roestelia</i> Rebent., <i>Prodr. fl. neomarch.</i> : 350. 1804. Type: <i>R. cancellata</i> (Pers.) Rebent. 1804, basionym: <i>Aecidium cancellatum</i> Pers. 1792, now regarded as <i>Gymnosporangium sabinae</i> (Dicks.) G. Winter 1884.	Protection needed by NCF.
Helicobasidium Pat. in <i>Bull. Soc. Bot. France</i> 32 : 172. 1885. Type: <i>H. purpureum</i> (Tul.) Pat. 1885, basionym: <i>Hypochnus purpureus</i> Tul. 1865.	<i>Thanatophytum</i> Nees, <i>Syst. Pilze</i> : 148. 1816. Type: <i>T. crocorum</i> (Pers.) Nees 1816, basionym: <i>Sclerotium crocorum</i> Pers. 1801, now regarded as <i>Helicobasidium purpureum</i> (Tul.) Pat. 1885. <i>Tuberculina</i> Tode ex Sacc. in <i>Michelia</i> 2 : 34. 1881. Type: <i>T. persicina</i> (Ditmar) Sacc. 1881, basionym: <i>Tubercularia persicina</i> Ditmar 1817, now regarded as <i>Helicobasidium purpureum</i> (Tul.) Pat. 1885.	Protection needed by NCF.
Hyalopycnis Höhn. in <i>Hedwigia</i> 60 : 152. 1918. Type: <i>H. hyalina</i> Höhn. 1918, now regarded as <i>Heterogastridium pycnidioideum</i> Oberw. & R. Bauer 1990.	<i>Heterogastridium</i> Oberw. & R. Bauer in <i>Mycologia</i> 82 : 57. 1990. Type: <i>H. pycnidioideum</i> Oberw. & R. Bauer 1990.	None.

Table 1. (Continued).

Recommended generic name	Synonymous alternate morph generic name(s)	Action required
Kriegeria Bres. in Rev. Mycol. (Toulouse) 13 : 14. 1891. Type: <i>K. eriophori</i> Bres. 1891.	<i>Xenogloea</i> Syd. & P. Syd. in Ann. Mycol. 17 : 44. 1919. Type: <i>X. eriophori</i> (Bres.) Syd. & P. Syd. 1919, basionym: <i>Kriegeria eriophori</i> Bres. 1891. <i>Zymoxenogloea</i> D.J. McLaughlin & Doublés in Mycologia 84 : 671. 1992. Type: <i>Z. eriophori</i> D.J. McLaughlin & Doublés 1992, now regarded as <i>Kriegeria eriophori</i> Bres. 1891.	None.
Kuehneola Magnus in Bot. Centralbl. 74 : 169. 1898. Type: <i>K. albida</i> (J.G. Kühn) Magnus 1898, basionym: <i>Chrysomyxa albida</i> J.G. Kühn 1883, now regarded as <i>Kuehneola uredinis</i> (Link) Arthur 1906.	<i>Spirechina</i> Arthur in J. Mycol. 13 : 30. 1907. Type: <i>S. loeseneriana</i> (Henn.) Arthur 1907, basionym: <i>Uredo loeseneriana</i> Henn. 1898, now regarded as <i>Kuehneola loeseneriana</i> (Henn.) H.S. Jacks. & Holw. 1931.	None.
Melampsorella J. Schröt. in Hedwigia 13 : 85. 1874. Type: <i>M. caryophyllacearum</i> (DC.) J. Schröt. 1874, basionym: <i>Uredo caryophyllacearum</i> DC. 1805.	<i>Peridermium</i> (Link) J.C. Schmidt & Kunze in Deutschl. Schwämme 6 : 4. 1817, basionym: <i>Hypodermium</i> subgen. <i>Peridermium</i> Link 1816. Type: <i>Aecidium elatinum</i> Alb. & Schwein. 1805, now regarded as <i>Melampsorella caryophyllacearum</i> (DC.) J. Schröt. 1874.	Protection needed by NCF.
Milesina Magnus in Ber. Deutsch. Bot. Ges. 27 : 325. 1909. Type: <i>M. kriegeriana</i> (Magnus) Magnus 1909, basionym: <i>Melampsorella kriegeriana</i> Magnus 1901.	<i>Milesia</i> F.B. White in Scott. Nat. 4 : 162. 1878. Type: <i>M. polypodii</i> F.B. White 1878, now regarded as <i>Milesina polypodii</i> (F.B. White) Aime & Rossman 2018.	Protection needed by NCF.
Phragmidium Link in Ges. Naturf. Freunde Berlin Mag. 7 : 30. 1816. Type: <i>P. mucronatum</i> (Pers.) Schltld. 1824, basionym: <i>Puccinia mucronata</i> Pers. 1794.	<i>Aregma</i> Fr., Observ. Mycol. 1 : 22., 1815. Type: <i>A. obtusatum</i> Fr. 1832, now regarded as <i>Phragmidium fragariae</i> (DC.) G. Winter 1884, basionym: <i>Puccinia fragariae</i> DC., 1808. <i>Lecythea</i> Lév. in Ann. Sci. Nat., Bot., sér. 3 8 : 373. 1847. Type: <i>L. miniata</i> (Pers.) Lév. 1847, basionym: <i>Uredo miniata</i> Pers. 1801, now regarded as <i>Phragmidium mucronata</i> (Pers.) Schltld. 1824.	Protection needed by NCF.
Prospodium Arthur in J. Mycol. 13 : 31. 1907. Type: <i>P. appendiculatum</i> (G. Winter) Arthur 1907, basionym: <i>Dicaeoma appendiculata</i> Kuntze 1898.	<i>Canasta</i> A.A. Carvalho & J.F. Hennen in Mycologia 102 : 1096. 2010. Type: <i>C. cruscula</i> A.A. Carvalho & J.F. Hennen 2010, now regarded as <i>Prospodium cruscula</i> (A.A. Carvalho & J.F. Hennen) Aime & Rossman 2018.	None.
Puccinia Pers. in Neues Mag. Bot. 1 : 118. 1794. Type: <i>P. graminis</i> Pers. 1794.	<i>Aecidium</i> Pers., Observ. Mycol. 1 : 97. 1796. Type: <i>A. berberidis</i> Pers. 1796, now regarded as <i>Puccinia graminis</i> Pers. 1794. <i>Caeoma</i> Link in Ges. Naturf. Freunde Berlin Mag. 3 : 5. 1809. Type: <i>C. berberidis</i> (Pers.) Har. 1809, basionym: <i>Aecidium berberidis</i> Pers. 1792, now regarded as <i>Puccinia graminis</i> Pers. 1794.	None.
Rhodotorula F.C. Harrison in Proc. & Trans. Roy. Soc. Canada, ser. 3 21 : 349. 1927. Type: <i>R. glutinis</i> (Fresen.) F.C. Harrison 1928.	<i>Chromotorula</i> F.C. Harrison in Proc. & Trans. Roy. Soc. Canada, ser. 3 21 : 350. 1927. Type: <i>C. kitae</i> F.C. Harrison 1927. <i>Rhodospordium</i> Banno in J. Gen. Appl. Microbiol. 13 : 192. 1967. Type: <i>R. toruloides</i> Banno 1967, now regarded as <i>Rhodotorula toruloides</i> (Banno) Q.M. Wang et al. 2015.	None.
Septobasidium Pat. in J. Bot. (Morot) 6 : 63. 1892. Type: <i>S. velutinum</i> Pat. 1892.	<i>Johncouchia</i> S. Hughes & Cavalc. in Canad. J. Bot. 61 : 2226. 1983. Type: <i>J. mangiferae</i> (Bat.) S. Hughes & Cavalc. 1983, basionym: <i>Cladotrichum mangiferae</i> Bat. 1956, now regarded as <i>Septobasidium pilosum</i> Boedijn & B.A. Steinm. 1930.	None.

Table 1. (Continued).

Recommended generic name	Synonymous alternate morph generic name(s)	Action required
<p><i>Sporobolomyces</i> Kluver & C.B. Niel in Centralbl. Bakteriol., 2 Abt. 2: 63. 1924.</p> <p>Type: <i>S. salmonicolor</i> (B. Fisch. & Brebeck) Kluver & C.B. Niel 1924, basionym: <i>Blastoderma salmonicolor</i> B. Fisch. & Brebeck 1894.</p>	<p><i>Aessosporon</i> Van der Walt in Antonie van Leeuwenhoek 36: 54. 1970.</p> <p>Type: <i>A. salmonicolor</i> Van der Walt 1970, now regarded as <i>Sporobolomyces salmonicolor</i> (B. Fisch. & Brebeck) Kluver & C.B. Niel 1924.</p> <p><i>Blastoderma</i> B. Fisch. & Brebeck, in Fischer, Morph. Syst.: 47. 1894.</p> <p>Type: <i>B. salmonicolor</i> B. Fisch. & Brebeck 1894, now regarded as <i>Sporobolomyces salmonicolor</i> (B. Fisch. & Brebeck) Kluver & C.B. Niel 1924.</p> <p><i>Rhodomycetes</i> Wettst. in Sitzungsber. Kaiserl. Akad. Wiss., Wien, Math.-naturw. Kl., Abt. 1 91: 39. 1885.</p> <p>Type: <i>R. kochii</i> Wettst. 1885, now regarded as <i>Sporobolomyces salmonicolor</i> (B. Fisch. & Brebeck) Kluver & C.B. Niel 1924.</p> <p><i>Sporidiobolus</i> Nyland in Mycologia 41: 686. 1950.</p> <p>Type: <i>S. johnsonii</i> Nyland 1950, now regarded as <i>Sporobolomyces johnsonii</i> (Nyland) Q.M. Wang <i>et al.</i> 2015.</p>	Protection needed by NCF.
<p><i>Uromyces</i> (Link) Unger, Exanth. Pflanzen: 277. 1833 (nom. cons.), basionym: <i>Hypodermium</i> subgen. <i>Uromyces</i> Link 1816.</p> <p>Type: <i>U. appendiculata</i> (Pers.) Unger 1796, basionym: <i>Uredo appendiculatus</i> Pers., now regarded as: <i>Uromyces appendiculatus</i> (Pers.) Link 1816.</p>	<p><i>Uredo</i> Pers., Syn. Meth. Fung. 1: 214. 1801.</p> <p>Type: <i>U. betae</i> Pers. 1801, now regarded as <i>Uromyces beticola</i> (Belynyck) Boerema, Loer. & Hamers 1987.</p>	Protection needed by NCF.

clarified the terminology associated with them, thus these generic names are synonyms. Given that *Basidiopycnis* has priority and both generic names are monotypic and relatively obscure (GS *Basidiopycnis* = 37, *Basidiopycnides* = 5), we follow the principle of priority and recommend the use of *Basidiopycnis*.

Use *Blastospora* Dietel 1908 (S) rather than *Pelastoma* Salazar-Yepes *et al.* 2012 (A)

The generic name *Blastospora* typified by *B. smilacis* includes tropical heteroecious rust fungi as summarized by Salazar-Yepes *et al.* (2012) who described *Pelostoma* typified by *P. juruense* for the asexual morph of *B. colombiana* (syn. *Hemileia colombiana*). Aime (2006) demonstrated that *B. smilacis* has affinities to *Hemileia* and *Maravalia* in the *Mikronegeriaceae*. *Blastospora smilacis* and *B. itoana* were considered to have asexual morphs belonging to *Caeoma* (Ono *et al.* 1986, 1987); however, the type of *Caeoma* is *C. berberidis*, a synonym of *Puccinia graminis*, thus *Caeoma* is a synonym of *Puccinia*. No other species of *Blastospora* nor the second species of *Pelostoma* have been sequenced. Assuming that *B. colombiana* is congeneric with *B. smilacis*, then *Blastospora* and *Pelostoma* are synonyms. At present five species are included in *Blastospora* and one name additional to the type has been placed in *Pelostoma*. Considering the fact that *Blastospora* has priority, a greater number of names, and is more commonly used (GS *Blastospora* = 567, *Pelostoma* = 11), we recommend the use of *Blastospora* for this genus.

Two new combinations are required:

***Blastospora juruensis* (Henn.) Aime & Rossman, comb. nov.**

MycoBank MB824643

Basionym: *Uredo juruensis* Henn., *Hedwigia* 43: 161 (1904).

Synonyms: *Hemileia juruensis* (Henn.) P. Syd. & Syd., *Mon. Ured.* 3: 219 (1915).

Hemileia jahnii Syd., *Ann. Mycol.* 28: 46 (1930).

Uredo jahnii (Syd.) Ritschel, *Biblioth. Mycol.* 200: 98 (2005).

Hemileia colombiana Buriticá, *Caldasia* 12: 165 (1978).

Blastospora colombiana (Buriticá) Salazar-Yepes *et al.*, *Mycologia* 104: 116 (2012).

***Blastospora subneurophylla* (Speg.) Aime & Rossman, comb. nov.**

MycoBank MB824644

Basionym: *Uredo subneurophylla* Speg., *Anales Soc. Ci. Argent.* 17: 123 (1883).

Synonyms: *Puccinia subneurophylla* (Speg.) Speg., *Anales Mus. Nac. Hist. Nat. Buenos Aires* 31: 384 (1922).

Pelastoma subneurophyllum (Speg.) Salazar-Yepes *et al.*, *Mycologia* 104: 116 (2012).

Uredo forsteroniae Henn., *Hedwigia* 34: 99 (1895).

Use *Camptobasidium* Marvanová & Suberkr. 1990 (S) rather than *Crucella* Marvanová & Suberkr. 1990 (A)

Table 2. Recommended generic names of *Ustilaginomycotina* among sexual-asexual names that compete for use. The recommended name is in bold; see text for the rationale behind these recommendations. For each generic name this list provides the author, its date and place of publication, type species of the genus, its basionym, their dates of publication and the currently accepted name, if different. The action required is indicated in the last column such as approval by the Nomenclature Committee for Fungi (NCF) for those generic names that do not have priority and thus need protection.

Recommended generic name	Synonymous alternate morph generic name(s)	Action required
<i>Anthracoidea</i> Bref., Unters. Gesamtgeb. Mykol. 12 : 143. 1895. Type: <i>A. caricis</i> (Pers.) Bref. 1896, basionym: <i>Uredo caricis</i> Pers. 1801.	<i>Crotalia</i> Liro in Ann. Acad. Sci. Fenn., Ser. A 42 (1): 545, 273. 1938. Type: <i>C. cintractiae-fischeri</i> Liro 1939, now regarded as <i>Anthracoidea fischeri</i> (P. Karst.) Kukkonen 1963.	None.
<i>Doassansia</i> Cornu in Ann. Sci. Nat., Bot., sér. 6 15 : 285. 1883. Type: <i>D. alismatis</i> (Nees ex Fr.) Cornu 1883, basionym: <i>Sclerotium alismatis</i> Nees ex Fr. 1822.	<i>Savulescuella</i> Cif. in Bontea et al. [eds], <i>Omag. Savulescu</i> : 179. 1959. Type: <i>S. alismacearum</i> (Sacc.) Cif. 1959, basionym: <i>Cylindrosporium alismacearum</i> Sacc. 1880, now regarded as <i>Doassansia alismatis</i> (Nees ex Fr.) Cornu 1883.	None.
<i>Doassansiopsis</i> (Setch.) Dietel in Engler & Prantl, Nat. Pflanzenfam. 1 (1**): 15. 1897, basionym: <i>Doassansia</i> subgen. <i>Doassansiopsis</i> Setch. 1891. Type: <i>Doassansia deformans</i> Setch. 1892, now regarded as <i>Doassansiopsis deformans</i> (Setch.) Dietel 1897.	<i>Doassansiella</i> Zambett. in Rev. Mycol. (Paris) 35 : 164. 1970. Type: <i>D. aquatilis</i> (Peck) Zambett. 1970, basionym: <i>Ramularia aquatilis</i> Peck 1884, now regarded as <i>Doassansiopsis hydrophila</i> (A. Dietr.) Lavrov 1937.	None.
<i>Entyloma</i> de Bary in Bot. Zeitung (Berlin) 32 (7): 101, 1874. Type: <i>E. microsporum</i> J. Schröt. 1874. Although some have attributed <i>E. microsporum</i> to <i>Protomyces microsporus</i> F. Unger 1834, this name is a later homonym of <i>P. microsporus</i> Pass. 1833.	<i>Entylomella</i> Höhn. in Ann. Mycol. 22 : 191. 1924. Type: <i>E. ranunculi</i> (Bonord.) Höhn. 1916, basionym: <i>Fusidium ranunculi</i> Bonord. 1851, now regarded as <i>Entyloma ficariae</i> F. Thümen ex A.A. Fischer v. Waldheim 1877.	None.
<i>Mycosarcoma</i> Bref., Unters. Gesamtgeb. Mykol. 15 : 53. 1912. Type: <i>M. maydis</i> (DC.) Bref. 1912, basionym: <i>Uredo maydis</i> DC. 1815. OR <i>Ustilago</i> (Pers.) Roussel, Fl. Calvados, ed. 2 : 4. 1806, basionym: <i>Uredo</i> [unranked] <i>Ustilago</i> Pers. 1801. Type: <i>Ustilago hordei</i> (Pers.) Lagerh. 1889, basionym: <i>Uredo segetum a hordei</i> Pers. 1801, or with a newly conserved type proposed by Thines (2016): <i>Ustilago maydis</i> (DC.) Corda 1842, basionym: <i>Uredo maydis</i> DC. 1815.	<i>Pseudozyma</i> Bandoni in J. Linn. Soc., Bot. 91 : 38. 1985. Type: <i>P. prolifica</i> Bandoni 1985, now regarded as <i>Mycosarcoma maydis</i> (DC.) Bref. 1912 or <i>Ustilago maydis</i> (DC.) Corda 1842.	Awaiting decision of NCF on the proposal of Thines (2016) as to whether <i>Mycosarcoma</i> should be taken up.
<i>Thecaphora</i> Fingerh. in Linnaea 10 : 230. 1836. Type: <i>T. hyalina</i> Fingerh. 1836, now regarded as <i>Thecaphora seminis-convolvuli</i> (Desm.) S. Ito 1935.	<i>Angiosorus</i> Thirum. & M.J. O'Brien 1974 in <i>Sydowia</i> 26 : 201 (1974) [1972]. Type: <i>A. solani</i> Thirum. & M.J. O'Brien 1974, now regarded as <i>Thecaphora solani</i> (Thirum. & M.J. O'Brien) Mordue 1988. <i>Rhombiella</i> Liro in Ann. Acad. Sci. Fenn., Ser. A 42 (1): 507, 545. 1939. Type: <i>R. cardamines</i> Liro 1939, now regarded as <i>Thecaphora thlaspeos</i> (Beck) Vánky 2004. <i>Thecaphorella</i> H. Scholz & I. Scholz in <i>Nova Hedwigia</i> 47 : 71. 1988. Type: <i>T. antherarum</i> (Oudem.) H. Scholz & I. Scholz 1988, basionym: <i>Gloeosporium antherarum</i> Oudem. 1898, now regarded as <i>Thecaphora seminis-convolvuli</i> (Desm.) S. Ito 1935.	None.
<i>Tilletia</i> Tul. & C. Tul. in Ann. Sci. Nat., Bot., sér. 3 7 : 112. 1847. Type: <i>Tilletia caries</i> (DC.) Tul. & C. Tul. 1847, basionym: <i>Uredo caries</i> DC. 1815, now regarded as <i>T. tritici</i> (Bjerk.) G. Winter 1875, basionym: <i>Lycoperdon tritici</i> Bjerk. 1775.	<i>Tilletiella</i> Zambett. in Rev. Mycol. (Paris) 35 : 165. 1970. Type: <i>T. alopecuri</i> (Sawada) Zambett. 1970, basionym: <i>Entyloma alopecuri</i> Sawada 1922, now regarded as <i>Tilletia alopecuri</i> (Sawada) L. Ling 1949.	None.

Table 2. (Continued).

Recommended generic name	Synonymous alternate morph generic name(s)	Action required
<i>Urocystis</i> Rabenh. ex Fuckel in Jahrb. Nassauischen Vereins Naturk. 23–24 : 41. 1870 (nom. cons.). Type: <i>U. occulta</i> (Wallr.) A.A. Fisch. Waldh. 1867, basionym: <i>Erysibe occulta</i> Wallr. 1833.	<i>Paepalopsis</i> J.G. Kühn in Irmischia 7 : 39, 1882. Type: <i>P. irmischiae</i> J.G. Kühn 1882, now regarded as <i>Urocystis primulae</i> (Rostr.) Vánky 1985. <i>Polycystis</i> Lév. in Ann. Sci. Nat. Bot., Ser. 3, 5 : 269. 1846. Type: <i>P. pompholygodes</i> Lév. 1846, now regarded as <i>Urocystis anemones</i> (Pers.) G. Winter 1880.	None.

The generic names *Camptobasidium* and *Crucella* were described in the same article based on sexually and asexually typified morphs of the same species, thus they are synonyms and compete equally for use (Marvanová & Suberkropp 1990). Both are monotypic and have been used to the same extent, although *Crucella* is also the name of a fossil radiolarian, thus it is difficult to determine its usage as a fungus. Given the confusion about the use of the name *Crucella* and because preference has previously been given to sexual morph names, we recommend the use of *Camptobasidium*.

Protect *Classicula* R. Bauer *et al.* 2003 (S) over *Naiadella* Marvanová & Bandoni 1987 (A)

The monotypic generic name *Classicula* is typified by *C. fluitans*, which was described as the sexual morph of *N. fluitans* (Bauer *et al.* 2003), the type species of the monotypic generic name *Naiadella*, thus these names are synonyms. Neither name is frequently used (GS *Classicula* = 73; *Naiadella* = 86); however, the family *Classiculaceae*, order *Classiculales*, and class *Classiculomycetes* are based on the name *Classicula*, thus we recommend the use of *Classicula*.

Use *Cronartium* Fr. 1815 (S) rather than *Endocronartium* Y. Hirats. 1969 (S)

The generic name *Cronartium* is well known for species that cause stem rusts on conifers in temperate regions, often with alternate hosts on herbaceous plants. One of the most notorious species is *C. flaccidum*, cause of Scots pine blister rust, of which *C. asclepiadeum*, the type species of *Cronartium*, is a synonym (Mordue & Gibson 1978). *Endocronartium* was established for species genetically similar to *Cronartium* except for being autoecious (i.e. completing their life-cycle on a single host) and endocyclic (lacking aecia and uredinia and having aecioid telia) (Hiratsuka 1969). The type species of *Endocronartium*, *E. harknessii*, cause of western gall rust, has long been known to be indistinguishable from species of *Cronartium* except for the life-cycle reductions (Epstein & Buurlage 1988). With the change in the ICN, these generic names are synonyms and the six taxa of *Endocronartium* should be recognized in the older, commonly used genus *Cronartium*.

All species of *Endocronartium* have been shown to belong in the genus *Cronartium* either as synonyms or distinct species within that genus. Several new combinations are made below. In addition, *Endocronartium pini* (Willd.) Y. Hirats. (syn. *Peridermium pini* (Willd.) Lév.) has been shown

to be a synonym of *C. flaccidum* (Hantuka *et al.* 2002).

Although some of the asexual morphs of species of *Cronartium* have been recognized in *Peridermium*, that generic name is now regarded as a synonym of *Melampsorella* based on the type species, *Aecidium elatinum*, which is a later name for *M. caryophyllacearum* (see below under *Melampsorella*). Thus, names in *Peridermium* must be recognized in the sexually typified genus in which the species belong, primarily *Cronartium* but also *Chrysomyxa* and *Coleosporium*.

The following new combinations are required:

Cronartium bethelii (Hedgc. & Long) Aime & Rossman, **comb. nov.**

MycoBank MB824645

Basionym: *Peridermium bethelii* Hedgc. & Long, *Phytopathology* **3**: 251 (1913).

Vogler & Bruns (1998) demonstrated that *Peridermium bethelii* belongs in *Cronartium*, as sister to *C. comandrae*. With the change in the ICN, it is possible to place this name in *Cronartium*.

Cronartium kurilense (Dietel) Y. Ono, **comb. nov.**

MycoBank MB824647

Basionym: *Peridermium kurilense* Dietel, *Bot. Jahrb. Syst.* **37**: 107 (1905).

Synonym: *Endocronartium sahoanum* var. *hokkaidoense* Imazu & Kakish., *Trans. Mycol. Soc. Japan* **33**: 174 (1992).

A blister rust fungus on *Pinus pumila* in the North Kuril Islands (southeast of Kamchatka, Russia) was described as *Peridermium kurilense*, distinct from *C. ribicola* (Dietel 1905). It was subsequently synonymized under *C. kamtschaticum* (Jørstad 1934, Hiratsuka *et al.* 1992). An endocyclic fungus on *Pinus pumila* distributed in northern Honshu, Japan, was named as *E. sahoanum* (Imazu *et al.* 1989), and a similar fungus on the same host species distributed in Hokkaido was separated as a variety, *E. sahoanum* var. *hokkaidoense*, distinct from *E. sahoanum* var. *sahoanum* on the basis of differences in spore and germ tube morphology (Imazu & Kakishima 1992). Based on the similarity in spore shape, size, surface structure, and PCR-RFLP pattern, *P. kurilense* and *E. sahoanum* var. *hokkaidoense* have been shown to be conspecific and distinct from *C. kamtschaticum*, which is now considered a synonym of *C. ribicola* (Imazu *et al.* 2000, Kim *et al.* 2010). Thus, *P. kurilense* must be placed in *Cronartium*.

Cronartium sahoanum (Imazu & Kakish.) Aime & Rossman, **comb. nov.**

MycoBank MB824648

Basionym: Endocronartium sahoanum Imazu & Kakish. *Trans. Mycol. Soc. Japan* 30: 308 (1989).

Cronartium yamabense (Saho & I. Takah.) Aime & Rossman, **comb. nov.**

MycoBank MB824649

Basionym: Peridermium yamabense Saho & I. Takah., *Trans. Mycol. Soc. Japan* 22: 33 (1981).

Synonym: Endocronartium yamabense (Saho & I. Takah.) Paclt, *Mitt. Deutsch. Dendrol. Ges.* 77: 227 (1987).

Use *Elateraecium* Thirum. et al. 1966 (A) rather than *Hiratsukamyces* Thirum. et al. 1975 (S)

The generic name *Hiratsukamyces*, typified by *H. salaciicola*, was established for the sexual morph of *Elateraecium salaciicola*, type species of *Elateraecium* (Thirumalachar et al. 1966, 1975), thus these generic names are synonyms. Two names were included in *Hiratsukamyces* when it was described, each also having a name in *Elateraecium* for the asexual morph. Since then two more names have been added to *Elateraecium* by Gjaerum & Reid (1983). Both generic names are obscure but given the greater number of names, priority, and slightly greater use (GS *Elateraecium* = 12, *Hiratsukamyces* = 11), we recommend the use of *Elateraecium*.

Use *Gerwasia* Racib. 1909 (S) rather than *Morispora* Salazar-Yepes et al. 2007 (A)

Gerwasia, typified by *G. rubi*, is a well-known genus of autoecious rust fungi that occurs on *Rosaceae*, primarily in the Asian and American tropics (Cummins & Hiratsuka 2003). Twenty-two species have been described in *Gerwasia* of which 12 have asexual morphs that have been placed in *Morispora*, typified by *M. tenella*, a species with a sexual morph regarded as *G. tenella* (Salazar-Yepes et al. 2007). Thus, assuming that *G. rubi* and *G. tenella* are congeneric as indicated by Salazar-Yepes et al. (2007) but without molecular data, *Gerwasia* and *Morispora* are synonyms. All names in *Morispora* already have a name in *Gerwasia*. Given its priority, greater number of names, and widespread use (GS *Gerwasia* = 112, *Morispora* = 6), the generic name *Gerwasia* is recommended for use.

Protect *Gymnosporangium* R. Hedw. ex DC. 1805 (S) over *Roestelia* Rebent. 1804 (A)

The genus *Gymnosporangium*, based on *G. fuscum*, now recognized as *G. sabinae*, includes ubiquitous rust fungi that have been well-studied (Yun et al. 2009). Index Fungorum lists 145 names in this genus. The type species of *Roestelia*, *R. cancellata*, is a synonym of *G. sabinae* (Alexopoulos 1940, Kern 1973), thus these generic names are synonyms. Although 55 names have been placed in *Roestelia*, their sexual morphs belong primarily in *Gymnosporangium*. The need to protect *Gymnosporangium* was recognized many years ago; however, the proposal to conserve *Gymnosporangium* over *Roestelia* was rejected for lack of adequate justification (Rogers 1953). Because *Gymnosporangium* is by far the most commonly used name rather than *Roestelia* (GS *Gymnosporangium* = 6050,

Roestelia = 598) and includes a greater number of species, we recommend the protection of *Gymnosporangium*.

Protect *Helicobasidium* Pat. 1885 (S) over *Thanatophytum* Nees 1816 (A) and *Tuberculina* Tode ex Sacc. 1880 (A)

Helicobasidium, typified by *H. purpureum*, includes 15 species, some of which cause plant diseases, such as *H. purpureum*, the cause of cosmopolitan violet root rot, and the closely related *H. mompa*, also causing violet root rot in Africa and Asia (Sayama et al. 1994). The asexual morph of *H. purpureum* has been regarded as *Thanatophytum crocorum* (Roberts 1999). Two species were mentioned in the protolog for *Thanatophytum*; Donk (1962) lectotypified this genus with *T. crocorum*. The generic name *Tuberculina* is lecto-typified by *T. persicina* (Clements & Shear 1931), a parasite of rust fungi. The relationship between these names all representing the same species was confirmed by Lutz et al. (2004) in which the life strategy of this unusual fungus was clarified. More specifically, they determined that *T. persicina* is a synonym of *Helicobasidium purpureum*. The commonly used name *H. purpureum* based on *Hypochnus purpureus* is herein proposed for protection over the older names *Sclerotium crocorum* and *Tubercularia persicina*. Among these three generic names, *Helicobasidium* is most commonly used (GS *Helicobasidium* = 2090, *Thanatophytum* = 67, *Tuberculina* = 487). Given the widespread use of *Helicobasidium* and the number of species in this genus, we recommend *Helicobasidium* for protection. In addition we recommend the name of the type species, *H. purpureum*, for protection.

Name for protection:

Helicobasidium purpureum (Tul.) Pat., *Bull. Soc. Bot. France* 32: 171 (1885).

Basionym: Hypochnus purpureus Tul., *Ann. Sci. Nat., Bot.*, sér. 5 4: 295 (1865); nom. prot. prop.

Rejected names and synonyms: Tuber parasiticum Bull., *Hist. champ. France* 1: 81 (1791).

Sclerotium crocorum Pers., *Syn. meth. fung.* 1: 119 (1801).

Rhizoctonia crocorum (Pers.) DC., in De Candolle & Lamarck, *Fl. franç.*, 3rd edn 5: 110 (1815).

Thanatophytum crocorum (Pers.) Nees, *Syst. Pilze*: 148 (1816) ["1816-17"].

Rhizoctonia medicaginis DC., in De Candolle & Lamarck, *Fl. franç.*, 3rd edn 5: 111 (1815).

Tubercularia persicina Ditmar, in Sturm, *Deutschl. Fl.*, Abt. 3 1: 99 (1817).

Tuberculina persicina (Ditmar) Sacc., *Fungi italice autogr. del.* 17-28: tab. 964 (1881).

Protonema brebissonii Desm., *Pl. crypt. N. France*: no. 651 (1834).

Helicobasidium brebissonii (Desm.) Donk, *Taxon* 7: 164 (1958).

Rhizoctonia rubiae M.J. Decne., *Rech. Anat. Physiol. Garance*: 55 (1837).

Rhizoctonia violacea Tul. & C. Tul., *Fung. hypog.*: 188 (1851).

Among the extensive list of synonymous names, *Helicobasidium purpureum* is the name in *Helicobasidium* most commonly used for this important and ubiquitous

plant pathogen (GS *Helicosporium purpureum* = 627, *H. berbissonii* = 7). Although several names for this fungus have been placed in *Rhizoctonia*, the type species of *Rhizoctonia* is *R. solani*, which represents a genus in *Agaricomycotina*.

Use *Hyalopycnis* Höhn. 1918 (A) rather than *Heterogastridium* Oberw. & R. Bauer 1990 (S)

The generic name *Hyalopycnis* is typified by *H. hyalina*. *Hyalopycnis blepharistoma*, a synonym of *H. hyalina*, was regarded as the asexual morph of *Heterogastridium pycnidoidium*, the type of the monotypic generic name *Heterogastridium* as explained by Oberwinkler *et al.* (1990) in describing the order *Heterogastridiales*, now placed in *Microbotryomycetes*, *Pucciniomycotina* (Aime *et al.* 2006, Toome & Aime 2014). Thus, the generic names *Heterogastridium* and *Hyalopycnis* are synonyms. Although the generic name *Heterogastridium* has been used slightly more frequently (GS *Heterogastridium* = 54, *Hyalopycnis* = 43) and serves as the basis for the ordinal name, *Hyalopycnis* has priority and its use circumvents the need for a name change for the only species, thus we recommend the use of *Hyalopycnis*. Its use does not affect the ordinal name.

Use *Kriegeria* Bres. 1891 (S) rather than *Xenogloea* Syd. & P. Syd. 1919 (S) and *Zymoxenogloea* D.J. McLaughlin & Doublés 1992 (A)

The monotypic generic name *Kriegeria* Bres., typified by *K. eriophori*, was determined to be legitimate by Toome *et al.* (2015). Previously it has been confused with *Kriegeria* Höhnelt 1914 and *Kriegeria* Rabenh. ex Seaver 1943, whose species are now placed in other genera in the ascomycete families *Helotiaceae* and *Rutstroemiaceae*. *Kriegeria* Bres. 1891 has also been confused with the later monotypic generic name *Xenogloea* typified by the same species, *X. eriophori*. An asexual morph of *K. eriophori*, as *X. eriophori*, was named *Zymoxenogloea eriophori*, monotype species of *Zymoxenogloea*, thus *Kriegeria* Bres. 1891, *Xenogloea*, and *Zymoxenogloea* are synonyms. Following Toome *et al.* (2015) and priority, we recommend the use of *Kriegeria*.

Use *Kuehneola* Magnus 1899 (S) rather than *Spirechina* Arthur 1907 (A)

The generic name *Kuehneola*, typified by *K. albida*, now regarded as *K. uredinis* (Ono 2015), includes about 25 species of rust fungi that occur primarily on species of *Rubus* and other *Rosaceae* (Cummins & Hiratsuka 2003). The generic name *Spirechina*, typified by *S. loeseneriana*, has been used for the asexual morph of species of *Kuehneola* and *Gerwasia*. *Spirechina loeseneriana* is regarded as the asexual morph of *K. loeseneriana* (Jackson 1931). Although 13 species had been named in *Spirechina*, all have been placed in *Gerwasia*, *Kuehneola*, or *Mainsia* (Kirk 2017). Given its priority, greater number of names, and more frequent use (GS *Kuehneola* = 590, *Spirechina* = 42), we recommend *Kuehneola* for use.

Protect *Melampsorella* J. Schröt. 1874 (S) over *Peridermium* (Link) J.C. Schmidt & Kunze 1817 (A)

The type species of *Melampsorella* is *M. caryophyllacearum*, while *Peridermium* is typified with the conserved type, *Aecidium elatinum*, as formally proposed by Hiratsuka (1974) and approved (Petersen 1975). Because the latter is regarded as a synonym of *M. caryophyllacearum* (Hiratsuka 1974), these generic names are synonyms. *Melampsorella* has five names in Index Fungorum. The asexually typified generic name *Peridermium* includes many species most of which have sexual morphs in *Cronartium* (Vogler & Bruns 1998). Given the relationship of species of *Peridermium* with both *Melampsorella* and *Cronartium*, we recommend abandoning the name *Peridermium* in favour of *Melampsorella*.

Although the type species of *Peridermium* dictates that this generic name is a synonym of *Melampsorella*, a number of names introduced in *Peridermium* belong in other rust genera, such as *Cronartium*, *Coleosporium* and several genera of *Pucciniastraceae*. An effort will be made to determine the correct generic placement of species described in *Peridermium*. Furthermore, an asexual morph can still be referred to as peridermium-like, using the term in a non-latinized form to describe a morphology.

Protect *Milesia* Magnus 1909 (S) over *Milesia* F.B. White 1878 (A)

The genus *Milesia* includes over 60 species, primarily occurring on *Abies* alternating with ferns in several families. The type species, *Milesia kriegeriana*, occurs on *Abies* and *Dryopteris* in Europe and is known to have an asexual morph recognized as *Milesia kriegeriana*. The type species of *Milesia* is *Milesia polypodii*, recognized as the uredinial state of *Milesia dieteliana* (Laundon 1967), thus *Milesia* and *Milesia* are synonyms. Although *Milesia* is the older generic name and has about the same number of names as *Milesia* (72 vs. 67, respectively), *Milesia* is in broader use especially in recent literature (e.g. Berndt 2007, Kakishima *et al.* 2016, Klenke & Scholler 2015, Smith 2015, Tanese & Negrean 2007, Wingfield *et al.* 2004). For these reasons, we recommend the protection and use of *Milesia*.

One new combination is required:

Milesia polypodii (F.B. White) Aime & Rossman, comb. nov.

MycoBank MB824650

Basionym: *Milesia polypodii* F.B. White, *Scott. Naturalist (Perth)*. 4: 163 (1878) ["1877-1878"]

Synonyms: *Milesia dieteliana* (Syd. & P. Syd.) Magnus, *Ber. Deutsch. Bot. Ges.* 27: 325 (1909).

Melampsorella dieteliana Syd. & P. Syd., *Ann. Mycol.* 1(6): 537 (1903).

Protect *Phragmidium* Link 1816 (S) over *Aregma* Fr. 1815 (S) and use rather than *Lecythea* Lév. 1847 (A)

The generic name *Phragmidium* is applied to a group of rust species that occur on *Rosaceae*. Recent molecular studies have shown that *Phragmidium* is a well-defined genus in *Phragmidiaceae* and that the type species of *Phragmidium*, *P. mucronatum*, is distinct from the other species commonly encountered on dog rose, *P. tuberculatum* (Ritz *et al.* 2005,

Yun et al. 2011). Both those papers also demonstrated that *P. mucronatum* is congeneric but distinct from *P. fragariae*, the correct name for the type species of *Aregma*, *A. obtusatum*; thus, *Aregma* and *Phragmidium* are synonyms. Several species are mentioned in the protolog for *Aregma*; this genus was lectotypified by Laundon (1965).

The asexually typified generic name *Lecythea* is typified by *L. miniata*, based on *Uredo miniata*, and now regarded as a synonym of *P. mucronatum*. Although *Lecythea* includes 27 names, all but four have been placed in other genera (see Index Fungorum). Most of the 16 names in *Aregma* have been placed or have synonyms in *Phragmidium*, while 250 names have been described in *Phragmidium*; this generic name is much more commonly used than *Aregma* (GS *Phragmidium* = 3770, *Aregma* = 244). Given its common usage and greater number of names, we recommend that *Phragmidium* be protected.

Use *Prospodium* Arthur 1907 (S) rather than *Canasta* A.A. Carvalho & J.F. Hennen 2010 (A)

The generic name *Prospodium* includes about 50 species occurring on *Bignoniaceae* and *Verbenaceae* in tropical regions (Cummins & Hiratsuka 2003, Hernández & Hennen 2003). The generic name *Canasta* was established for asexual morphs of species of *Prospodium* (Carvalho & Hennen 2010), although the type species *C. cruscula* does not have a known sexual morph. Given that the other two names described in *Canasta* have sexual morphs that are species of *Prospodium*, it seems likely that *Canasta* is a synonym of *Prospodium*. Given its priority, greater number of species and widespread use, we recommend the use of *Prospodium*. One new combination is necessary:

Prospodium crusculum (A.A. Carvalho & J.F. Hennen)

Aime & Rossman, **comb. nov.**

Mycobank MB824651

Basionym: *Canasta cruscula* A.A. Carvalho & J.F. Hennen, *Mycologia* **102**: 1098 (2010)

Use *Puccinia* Pers. 1794 (S) rather than *Aecidium* Pers. 1796 (A) and *Caeoma* Link 1809 (A)

The generic name *Puccinia* is typified by *P. graminis*, the cause of the important wheat rust disease that has plagued this major food source throughout history (Schumann 1991). Several thousand names have been placed in *Puccinia*. The asexually typified generic name *Aecidium* is typified by *A. berberidis*, a name used for one of the alternate morphs of *P. graminis* (Cummins 1971), thus these generic names are synonyms. Prior to 2012, *Aecidium* was used when a sexual morph was lacking for rust fungi that appeared to have affinities with *Puccinia*. Before the changes in the ICN, the species could not be described in the "correct" genus (*Puccinia*) even though it was possible to determine its generic placement. The need for the generic name *Aecidium* no longer exists. The type species of *Caeoma*, *C. berberidis*, is a synonym of *Puccinia graminis* (Laundon 1967), thus *Caeoma* is a synonym of *Puccinia*. Given the ubiquitous use of the generic name *Puccinia*, it is recommended for use.

Use *Rhodotorula* F.C. Harrison 1928 (A) rather than *Chromotorula* F.C. Harrison 1928 (A) and *Rhodospordium* Banno 1967 (S)

The type species of *Rhodotorula*, *R. glutinis*, was shown to be congeneric with *Rhodospordium toruloides*, type of *Rhodospordium*, by Wang et al. (2015a). They recognized the priority of *Rhodotorula* and placed *Rhodospordium toruloides* in *Rhodotorula*. All 14 names described in *Rhodospordium* have been placed in two genera, *Rhodospordiobolus* and *Rhodotorula*. More than 150 names have been introduced in *Rhodotorula*. *Chromotorula*, typified by *C. kitae*, was published in the same article as *Rhodotorula* and thus has equal priority to *Rhodotorula*. *Chromotorula* has long been considered a synonym of *Rhodotorula* (Statzell-Tallman & Fell 1998), and all but two of the six names in *Chromotorula* have been placed elsewhere. In agreement with Wang et al. (2015a) as well as priority, greater number of names, and most frequent use (GS *Rhodotorula* = 50 300, *Rhodospordium* = 6130 *Chromotorula* = 30), we recommend the use of *Rhodotorula*.

Use *Septobasidium* Pat. 1892 (S) rather than *Johncouchia* S. Hughes & Cavalc. 1983 (A)

The monotypic generic name *Johncouchia*, typified by *J. mangiferae*, was described for the asexual morph of *Septobasidium pilosum* (Hughes & Cavalcanti 1983). *Septobasidium pilosum* and *S. velutinum*, the type species of *Septobasidium*, were determined to be congeneric by Henk & Vilgalys (2007). *Septobasidium* includes over 300 names, has priority and is widely used (GS *Septobasidium* = 1200, *Johncouchia* = 15), and so we recommend the use of *Septobasidium*.

Septobasidium has already been formally conserved against *Campylobasidium* Lagerh. ex F. Ludw. 1892, *Gausapia* Fr. 1825, and *Glenospora* 1849.

Protect *Sporobolomyces* Kluyver & C.B. Niel 1924 (A) over *Blastoderma* B. Fisch. & Breback 1894 (A) and *Rhodomyces* Wettst. 1885 (A) and use rather than *Aessosporon* Van der Walt 1970 (S) and *Sporidiobolus* Nyland 1950 (S)

The generic name *Sporobolomyces*, typified by *S. salmonicolor*, includes about 100 names and is widely used. An earlier generic name, *Rhodomyces*, typified by *R. kochii*, is now considered a synonym of *S. salmonicolor* (Sampaio 2011). A second earlier monotypic generic name, *Blastoderma*, is typified by *B. salmonicolor*, the basionym of *S. salmonicolor*, thus these three generic names are synonyms. Another later name, *Sporidiobolus*, has six species, and is typified by *S. johnsonii*, a name now placed in *Sporobolomyces* (Wang et al. 2015a). The generic name *Aessosporon*, typified by *A. salmonicolor*, was described as the sexual morph of *Sporobolomyces salmonicolor* (Van der Walt 1970). Later authors showed that *A. salmonicolor* had the same heterothallic life-cycle and was conspecific with *S. salmonicolor* (Fell & Statzell-Tallman 1980), thus *Aessosporon* and *Sporobolomyces* are synonyms. Among these five generic names *Blastoderma* has rarely been used. Although the name *Rhodomyces* appears in the literature,

the three described species are not closely related to the type, as shown by Wang *et al.* (2015a). A second species of *Aessosporon*, *A. dendrophilum*, was described as the sexual morph of *Bullera dendrophilum* (van der Walt 1973), but is now regarded as *Kwoniella dendrophila* in *Agaricomycotina* (Liu *et al.* 2015), and unrelated to *Sporobolomyces*. Because *Sporobolomyces* has priority, includes about 100 names and is widely used (GS *Sporobolomyces* = 10 600 *Rhodomycetes* = 183, *Blastoderma* = 523, *Sporidiobolus* = 2970, *Aessosporon* = 171), we recommend that *Sporobolomyces* be protected over *Rhodomycetes* and *Blastoderma*, and that *Sporobolomyces* be used rather than *Sporidiobolus* and *Aessosporon*.

Protect *Uromyces* (Link) Unger 1833 (S) over *Uredo* Pers. 1801 (A)

The generic name *Uromyces*, typified by *U. appendiculatus*, includes over 800 names. When Persoon (1801) described *Uredo*, he included 30 species, one of which was *U. betae*. Laundon (1970) reviewed the history of the typification of the generic name *Uredo* and concluded that *U. betae* should be accepted as the type of the genus. *Uredo betae* is now considered a synonym of *Uromyces beticola* (Boerema *et al.* 1987, Simpson *et al.* 2006). *Uromyces appendiculatus* and *U. beticola* are regarded as congeneric, although species of *Uromyces* are interspersed within *Puccinia* in *Pucciniaceae* (Aime 2006, Demers *et al.* 2017, Maier *et al.* 2003, Wingfield *et al.* 2004), thus *Uromyces* and *Uredo* are synonyms. *Uredo* includes over 2000 names, but is not as widely cited as *Uromyces* (GS *Uromyces* = 26 400, *Uredo* 7500). Thus, we recommend that *Uromyces* be protected over *Uredo*.

The generic name *Uredo* has been used for uredinal morphs of rust fungi belonging to multiple rust families, and does not reflect the affinities of species. Prior to 2011, this generic name was used when a sexual morph was lacking and the species could not be described in the “correct” sexually typified genus according the *Code* even when it was possible to determine its generic placement. Now that placement of fungi in genera is no longer restricted by whether their types are asexual or sexual morphs, in theory the need for the generic name *Uredo* no longer exists. However, in reality rust fungi are often found without the sexual morph and it can be difficult to determine their affinities; the issue of genera typified by a sexual morphs is discussed further above (p. 76).

The generic name *Uromyces* is already conserved over *Coeomurus* Link ex Gray 1821 and *Pucciniola* L. Marchand 1829.

COMPETING SEXUAL AND ASEXUAL GENERIC NAMES OF *USTILAGINOMYCOTINA*, *BASIDIOMYCOTA* AND RECOMMENDATIONS FOR USE

Use *Anthracoidea* Bref. 1895 (S) rather than *Crotalia* Liro 1938 (A)

The sexually typified genus *Anthracoidea* in *Ustilaginales* includes over 200 names, of which 90 are accepted in Vánky (2012), while the asexually typified *Crotalia* has five names, all of which also have names in *Anthracoidea*. The type species of *Anthracoidea* is *A. caricis*. Initially the type

species of *Crotalia*, *C. cintractiae-fischeri*, was considered to be the asexual morph of *Cintractia fischeri*; however, that species is now regarded as *Anthracoidea fischeri* (Vánky 2012). Although *A. fischeri* was not included in Hendrichs *et al.* (2005), both *A. caricis* and *A. fischeri* are placed in *Anthracoidea* by Vánky (2012), thus *Anthracoidea* and *Crotalia* are synonyms.

Given its priority, the number of species, and more frequent citation in the literature (GS *Anthracoidea* = 619, *Crotalia* = 234), we recommend the use of *Anthracoidea*.

Use *Doassansia* Cornu 1883 (S) rather than *Savulescuella* Cif. 1959 (A)

The generic name *Doassansia*, typified by *D. alismatis*, includes about 12 species (Vánky 2012) while *Savulescuella*, typified by *S. alismacearum*, includes only three names. Vánky (2012) assumed that the asexual morph typified by *S. alismacearum* might be just “basidiospores of spores that germinated *in situ*”, nevertheless *Savulescuella* was validly published. Currently, *S. alismacearum* is considered a synonym of *D. alismatis*, making these names generic synonyms. Given its priority, number of species, and widespread use (GS *Doassansia* = 457, *Savulescuella* = 7), we recommend the use of *Doassansia*.

Use *Doassansiopsis* (Setch.) Dietel 1897 (S) rather than *Doassansiella* Zambett. 1970 (A)

The generic name *Doassansiopsis* is typified by *D. deformans*. Sixteen additional names have been placed in this genus, most of which are accepted by Vánky (2012). The monotypic generic name *Doassansiella* is typified by *D. aqualitis* and has rarely been used. The type species is now regarded as *Doassansiopsis hydrophila* (Vánky 2012). Given the priority, number of species, and more extensive use (GS *Doassansiopsis* = 153, *Doassansiella* = 2), we recommend the use of *Doassansiopsis*.

Doassansiopsis in *Urocystales* should not be confused with *Doassansia* and *Doassinga* in the *Doassansiales* (Begerow *et al.* 2004).

Use *Entyloma* de Bary 1874 (S) rather than *Entylomella* Höhn. 1924 (A)

The generic name *Entylomella* was first mentioned by Höhnel (1916), although in that publication a description is lacking of either the genus or species although one species, *E. ranunculi*, is listed. Later in discussing *Cylindrosporium*, Höhnel (1924) mentioned *Entylomella* again as the conidial form of *Entyloma* and *Doassansia* species and two pages later provided a description of the type species as discussed by Braun (1995). *Entylomella ranunculi* is the type of *Entylomella*, which is now regarded as a synonym of *Entyloma ficariae*. The type species of *Entyloma* is *E. microsporium*. Begerow *et al.* (2002) demonstrated that *E. ficariae* and *E. microsporium* belong in *Entyloma*, thus *Entylomella* and *Entyloma* are synonyms. Over 400 names are listed in *Entyloma* and this generic name is well known (GS = 2410) while 59 names have been described in *Entylomella*, many of which have been placed elsewhere, and the name has been used infrequently (GS = 60). Given the priority, greater number of names, and familiarity, we recommend the use of *Entyloma*.

Use *Mycosarcoma* Bref. 1912 or *Ustilago* (Pers.) Roussel 1806 (S) rather than *Pseudozyma* Bandoni 1985 (A)

The yeast-like morphs of *Ustilaginales* have been described in the genus *Pseudozyma*. Boekhout (1995) showed that the type species of *Pseudozyma*, *P. prolifica*, was closely related or synonymous with *Ustilago maydis*, the cause of corn smut (Ainsworth 1965). While 26 names have been described in *Pseudozyma*, many of these have now been placed in diverse sexual genera such as *Anthracoystis*, *Dirkmeia*, *Kalmanozyma*, *Langdonia*, *Moesziomyces* and *Sporisorium*. A controversy exists over which generic name to use for the species long referred to as *U. maydis*, an earlier name for *P. prolifica*. The generic name *Ustilago* had been considered to be typified by *U. hordei*, cause of barley smut (Clinton 1906, Vánky 2012). This and related species, such as *U. avenae*, *U. striiformis*, and *U. tritici* were determined by McTaggart et al. (2016) to fall in a clade distinct from the one that included *U. maydis*, type of the resurrected generic name *Mycosarcoma*, which was based on morphological and cultural characteristics by Brefeld (1912). Based on molecular data, five names were included in *Mycosarcoma* by McTaggart et al. (2016), resurrecting *Mycosarcoma maydis* for the corn smut and closely related species. However, a proposal has been made to conserve the name *Ustilago* with *U. maydis* as the conserved type (Thines 2016). *Pseudozyma prolifica* is a name used for the asexual morph of *Mycosarcoma / Ustilago maydis*, thus we recommend the use of *Mycosarcoma* or *Ustilago* depending on whether the NCF votes to conserve the name *Ustilago* with a new type. Because the generic name *Pseudozyma* is no longer available, the taxonomic treatment of several names in *Pseudozyma* (e.g. *P. alboarmeniaca*, *P. thailandica*, *P. tsukubaensis*, *P. pruni*) is unclear as they cannot currently be placed with confidence in any sexually typified lineage (Wang et al. 2015a; these species are therefore temporarily retained in *Pseudozyma* “pro tem.”

Use *Thecaphora* Fingerh. 1836 (S) rather than *Angiosorus* Thirum. & M.J. O'Brien 1974 (S), *Rhombiella* Liro 1939 (A) and *Thecaphorella* H. Scholz & I. Scholz 1988 (A)

The type species of *Thecaphora*, *T. hyalina*, is now regarded as *T. seminis-convolvuli* (Vánky et al. 2008). The type species of the asexually typified generic name *Rhombiella*, *R. cardamines*, is now regarded as *Thecaphora thlaspeos* (Vánky et al. 2008). The type of *Thecaphorella*, *T. antherarum*, was described as the asexual morph of *Thecaphora seminis-convolvuli* (Scholz & Scholz 1988). Begerow et al. (1997) demonstrated that *Thecaphora* represents a well-resolved genus in *Ustilaginales* including the types of *T. seminis-convolvuli* and *T. thlaspeos*; these three generic names are therefore synonyms. Similarly, Andrade et al. (2004) and Vánky et al. (2008) demonstrated that *Thecaphora* includes *T. seminis-convolvuli* and *T. solani* (syn. *Angiosorus solani*, type of the monotypic generic name *Angiosorus*), an economically important fungus causing potato smut (Mordue 1988). Vánky (2012) includes over 60 species in *Thecaphora*, under which he lists four sexually typified generic names as later synonyms, along with the rejected name *Sorosporium*. *Thecaphora* has

priority, includes numerous species, and is well known (GS *Thecaphora* = 1420, *Rhombiella* = 5, *Angiosorus* = 128), thus we recommend the use of *Thecaphora*.

Use *Tilletia* Tul. & C. Tul. 1847 (S) rather than *Tilletiella* Zambett. 1970 (A)

The genus *Tilletia* has long been accepted as the name for bunt fungi with over 100 accepted species, some of which cause serious diseases of crops in the grass family such as common bunt and dwarf bunt of wheat (Hoffman 1982, Mathre 1996). *Tilletia*, typified by *T. caries*, and treated as *T. tritici*, is now placed in *Tilletiaceae* (*Tilletiales*, *Exobasidiomycetes*; Begerow et al. 2006). The monotypic *Tilletiella* was established as an asexually typified genus for *T. alopecuri*, but that species is now regarded as *Tilletia alopecuri* (Vánky 2012). *Tilletiella* is therefore a synonym of *Tilletia*. Given the priority, large number of species, and widespread use (GS *Tilletia* = 16 900, *Tilletiella* = 0), we recommend the use of *Tilletia*.

Use *Urocystis* Rabenh. ex Fuckel 1870, nom. cons. (S) rather than *Paepalopsis* J.G. Kühn 1882 (A)

The type species of *Paepalopsis*, *P. irmischiae*, is considered the asexual morph of *Urocystis primulae*, a species included in *Urocystis*, which is typified by *U. occulta* (Vánky 2012); these generic names are consequently synonyms. Vánky (2012) listed four sexually typified generic synonyms of *Urocystis*, including *Tubercinia* Fr. 1832 nom. sanct. and *Polycystis* Lév., earlier names of *Urocystis* which has been conserved over them (Wakefield 1939). Three names have been placed in *Paepalopsis*, two of which are now regarded as species of *Urocystis*, while the third, *P. deformans*, is now placed in the ascomycete genus *Hapalosphaeria* as *H. deformans*. Vánky (2012) accepted over 160 species of *Urocystis* in his world monograph of smut fungi. Given the priority, number of species, and widespread use (GS *Urocystis* = 4920, *Paepalopsis* = 24), *Urocystis* remains the correct name for this genus.

ACKNOWLEDGEMENTS

We are extremely grateful to the following individuals for their input, discussions, and suggestions: Gaddam Bagyanarayana, Cvetomir Denchev, David McLaughlin, Alistair McTaggart, Drew Minnis, Franz Oberwinkler (†), Angela Schäfer, Roger Shivas, John Wiersema, and Andrey Yurkov.

REFERENCES

- Aime MC (2006) Toward resolving family-level relationships in rust fungi (*Uredinales*). *Mycoscience* **47**: 112–122.
- Aime MC, Matheny PB, Henk DA, Frieders EM, Nilsson RH, et al. (2006) An overview of the higher-level classification of *Pucciniomycotina* based on combined analyses of nuclear large and small subunit rDNA sequences. *Mycologia* **98**: 896–905.
- Ainsworth GC (1965) *Ustilago maydis*. *Descriptions of Pathogenic Fungi and Bacteria* **79**: 1–2.
- Alexopoulos CJ (1940) Some fungi from Greece. *Mycologia* **32**: 336–358.

- Andrade O, Moñoz G, Galdames R, Durán P, Honorato R (2004) Characterization, *in vitro* culture, and molecular analysis of *Thecaphora solani*, the causal agent of potato smut. *Phytopathology* **94**: 875–882.
- Bandoni RJ, Boekhout T (1998) *Agaricostilbum* Wright. In: *The Yeasts: a taxonomic study* (Kurtzman CP, Fell JW, eds): 639–640. 4th edn. Amsterdam: Elsevier.
- Bandoni R, Oberwinkler F (1981) *Hyalopycnis blepharistoma*: a pycnidial basidiomycete. *Canadian Journal of Botany* **59**: 1613–1620.
- Baral HO, Weber E, Gams W, Hagedorn G, Liu B *et al.* (2017) ["2018"] Generic names in the *Orbiliaceae* (*Orbiliomycetes*) and recommendations on which names should be protected or suppressed. *Mycological Progress* **17**: 5–31.
- Bauer R, Begerow D, Oberwinkler F, Marvanová L (2003) *Classicula*: the teleomorph of *Naiadella fluitans*. *Mycologia* **95**: 756–764.
- De Beer ZW, Seifert KA, Wingfield MJ (2013) A nomenclator for ophiostomatoid genera and species in the *Ophiostomatales* and *Microascales*. In: *Ophiostomatoid Fungi: expanding frontiers* (Seifert KA, De Beer ZW, Wingfield MJ, eds): 245–322. [CBS Biodiversity Series no. 12.] Utrecht: CBS-KNAW Fungal Biodiversity Centre.
- Begerow D, Bauer R, Oberwinkler F (1997) Phylogenetic studies on nuclear LSU rDNA sequences of smut fungi and related taxa. *Canadian Journal of Botany* **75**: 2045–2056.
- Begerow D, John B, Oberwinkler F (2004). Evolutionary relationships among β -tubulin gene sequences of basidiomycetous fungi. *Mycological Research* **108**: 1257–1263.
- Begerow D, Lutz M, Oberwinkler F (2002) Implications of molecular characters for the phylogeny of the genus *Entyloma*. *Mycological Research* **106**: 1392–1399.
- Begerow D, Stoll M, Bauer R (2006) A phylogenetic hypothesis of *Ustilaginomycotina* based on multiple gene analyses and morphological data. *Mycologia* **98**: 906–916.
- Berndt R (2015) The rust fungi (*Uredinales*) on ferns in South Africa. *Mycological Progress* **7**: 7–19.
- Boekhout T (1995) *Pseudozyma* Bandoni emend Boekhout, a genus for yeast-like anamorphs of *Ustilaginales*. *Journal for General Applied Microbiology* **41**: 359–366.
- Boerema GH, Loerakker WM, Hamers MEC (1987) Checklist for scientific names of common parasitic fungi. Supplement Series 2a (additions and corrections): Fungi on field crops: beet and potato; caraway, flax and oilseed poppy. *Netherlands Journal of Plant Pathology* **93** (Suppl.): 1–20.
- Braun U (1995) *A Monograph of Cercospora, Ramularia and Allied Genera (Phytopathogenic Hyphomycetes)*. Vol. 1. Eching: IHW-Verlag.
- Braun U (2013) (2210-2232) Proposals to conserve the teleomorph-typified name *Blumeria* against the anamorph-typified name *Oidium* and twenty-two teleomorph-typified powdery mildew species names against competing anamorph-typified names (*Ascomycota: Erysiphaceae*). *Taxon* **62**: 1328–1331.
- Brefeld O (1812) *Untersuchungen aus dem Gesamtgebiete der Mykologie*. Vol. 15. *Die Brandpilze und die Brandkrankheiten* **5**: 1–151. Münster: Commissions-Verlag v. H. Schöningh.
- Castlebury LA, Carris LM, Vánky K (2005) Phylogenetic analysis of *Tilletia* and allied genera in order *Tilletiales* (*Ustilaginomycetes; Exobasidiomycetidae*) based on large subunit nuclear rDNA sequences. *Mycologia* **97**: 888–900.
- Clements FE, Shear CL (1931) *The Genera of Fungi*. New York, NY: H.W. Wilson Co.
- Clinton GP (1906) Order *Ustilaginales*. *North American Flora* **7**: 1–82.
- Cummins GB (1971) *The Rust Fungi of Cereals, Grasses and Bamboos*. New York: Springer-Verlag.
- Cummins GB, Hiratsuka Y (2003) *Illustrated Genera of Rust Fungi*. 3rd edn. St Paul, MN: American Phytopathological Society Press.
- Daniel H-M, Lachance M-A, Kurtzman CP (2014) On the reclassification of species assigned to *Candida* and other ascomycetous yeast genera based on phylogenetic circumscription. *Antonie van Leeuwenhoek* **10**: 67–84.
- De Carvalho AA, Hennen JF (2010) New species and nomenclature in *Prospodium* (*Uropyxidaceae, Pucciniales*) and the new anamorphic genus *Canasta* in the Neotropics. *Mycologia* **102**: 1096–1113.
- Demers JE, Liu M, Hambleton S, Castlebury LA (2017) Rust fungi on *Panicum*. *Mycologia* **109**: 1–17.
- Dietel P (1905) *Uredinieae Japonicae*. VI. *Botanisches Jahrbücher* **37**: 97–109.
- Donk MA (1962) The generic names proposed for *Hymenomycetes*. XII, *Deuteromycetes*. *Taxon* **11**: 75–104.
- Epstein L, Buurlage MB (1988) Nuclear division in germinating aeciospores and its taxonomic significance for the western gall rust fungus, *Peridermium harknessii*. *Mycologia* **80**: 235–240.
- Fell JW (1966) *Sterigmatomyces*, a new fungal genus from marine areas. *Antonie van Leeuwenhoek* **32**: 99–104.
- Fell JW, Statzell-Tallman A (1980) Mating between strains of the yeasts *Aessosporon salmonicolor* and *Sporobolomyces* spp. *International Journal of Systematic Bacteriology* **30**: 206–207.
- Gjaerum HB, Reid DA (1983) Three new species and a new combination in *Uredinales*. *Transactions of the British Mycological Society* **81**: 650–654.
- Hantula J, Kasanen R, Kaitera J, Moricca S (2002) Analysis of genetic variation suggest that pine rusts *Cronartium flaccidum* and *Peridermium pini* belong to the same species. *Mycological Research* **106**: 203–209.
- Hausner G, Reid J, Eyjólfssdóttir GG, Iranpour M, Loewen P (2008) *Basidiopycnides albertensis* gen. et sp. nov., a new anamorphic fungus with phylogenetic affinities in the *Atractiellales* (*Basidiomycota*). *Mycotaxon* **103**: 279–297.
- Healy R, Pfister DH, Rossman AY, Marvanová L, Hansen K (2016) Competing sexual-aseexual generic names of *Pezizomycetes* and recommendations for use. *IMA Fungus* **7**: 285–288.
- Hendrichs M, Begerow D, Bauer R, Oberwinkler F (2005) The genus *Anthracoidea* (*Basidiomycota, Ustilaginales*): a molecular phylogenetic approach using LSU rDNA sequences. *Mycological Research* **109**: 31–40.
- Henk DA, Vilgalys R (2007) Molecular phylogeny suggests a single origin of insect symbiosis in the *Pucciniomycetes* with support for some relationships within the genus *Septobasidium*. *American Journal of Botany* **94**: 1515–1526.
- Hernández JR, Hennen JF (2003) Rust fungi causing galls, witches' brooms, and other abnormal plant growths in northwestern Argentina. *Mycologia* **95**: 728–755.
- Hiratsuka N, Sato S, Katsuya K, Kakishima M, Hiratsuka Y, *et al.* (1992) *The Rust Flora of Japan*. Tsukuba: Tsukuba.
- Hiratsuka Y (1969) *Endocronartium*, a new genus for autoecious pine stem rusts. *Canadian Journal Botany* **47**: 1493–1495.
- Hiratsuka Y (1974) (373) Proposal to conserve the generic name *Peridermium* (Link) Schmidt & Kunze with a conserved type species, *Aecidium elatinum* Alb. & Schw. (*Fungi imperfecti uredinearum*). *Taxon* **23**: 428–429.
- Hoffmann JA (1982) Bunt of wheat. *Plant Disease* **66**: 979–986.

- Höhnel F (1916) Mykologisches. *Österreich Botanische Zeitung* **66**: 94–112.
- Höhnel F (1924) Beitrag zur Kenntnis der Gattung *Cylindrosporium* Grev. *Annales Mycologici* **22**: 191–203.
- Hughes SL, Cavalcanti WA (1983) *Janetia mangiferae* n.sp. and the *Johncouchia* n. gen. anamorph of *Septobasidium pilosum* from Brasil. *Canadian Journal of Botany* **61**: 2224–2230.
- Imazu M, Azbukina ZM, Kakishima M, Fukushima K, Hishimura K, et al. (2000) Identification of a rust fungus on *Pinus pumila* collected in the North Kurils, Russia. *Mycoscience* **41**: 139–144.
- Imazu M, Kakishima M (1992) A new variety of *Endocronartium sahoanum* found on *Pinus pumila* in Hokkaido, Japan. *Transactions of the Mycological Society of Japan* **33**: 167–176.
- Imazu M, Kakishima M, Kaneko S (1989). *Endocronartium sahoanum*, a new stem rust fungus on *Pinus pumila* in Japan. *Transactions of the Mycological Society of Japan* **30**: 301–310.
- Jackson HS (1931) The rusts of South America based on the Holway collections -III. *Mycologia* **23**: 96–116.
- Johnston PR, Seifert KA, Stone JK, Rossman AY, Marvanová L (2014) Recommendations on generic names competing for use in *Leotiomycetes* (Ascomycota). *IMA Fungus* **5**: 91–120.
- Jørstad I (1934) A study on Kamtchatka *Uredinales*. *Skrifter utgitt av Det Norske Videnskaps-Akademi I Oslo, nat.-naturv. klasse* **1933**(9): 1–183.
- Kakishima M, Ji J-X, Wang Q, Li Y (2016) First report of rust disease caused by *Milesina dryopteridis* on two species of ferns, *Rumohra adiantiformis* and *Pteris fauriei*, in Japan. *Plant Disease* **100**: 2529.
- Kepler RM, Luangsa-ard JJ, Hywel-Jones NL, Quandt CA, Sung G-H, et al. (2017) A phylogenetic-based nomenclature for *Cordycipitaceae* (Hypocreales). *IMA Fungus* **8**: 335–354.
- Kern FD (1973) *A Revised Taxonomic Account of Gymnosporangium*. University Park, PN: Pennsylvania State University Press.
- Kim MS, Klopfenstein NB, Ota Y, Lee SK, Woo KS, et al. (2010) White pine blister rust in Korea, Japan and other Asian regions: comparisons and implications for North America. *Forest Pathology* **40**: 382–401.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA (2008) *Ainsworth & Bisby's Dictionary of the Fungi*. 10th edn. Wallingford: CAB International.
- Kirschner R, Begerow D, Oberwinkler, F (2001) A new *Chionosphaera* species associated with conifer inhabiting bark beetles. *Mycological Research* **105**: 1403–1408.
- Kirschner R, Oberwinkler F (2009) Supplementary notes on *Basidiopycnis hyalina* (Basidiomycota, Atractiellales) and its anamorph. *Mycotaxon* **109**: 29–38.
- Klenke F, Scholler M (2015) *Pflanzenparasitische Kleinpilze*. Berlin: Springer.
- Laundon GF (1965) The generic names of *Uredinales*. *Mycological Papers* **99**: 1–24.
- Laundon GF (1967) The taxonomy of the imperfect rusts. *Transactions of the British Mycological Society* **50**: 349–353.
- Laundon GF (1970) The lectotype of *Uredo*. *Taxon* **19**: 947.
- Liu XZ, Wang QM, Göker M, Groenwald M, Kachalkin AV et al. (2015) Towards an integrated phylogenetic classification of the *Tremellomycetes*. *Studies in Mycology* **81**: 85–147.
- Lutz M, Bauer R, Begerow D, Oberwinkler F (2004) *Tuberculina-Thanatophytum/Rhizoctonia crocorum-Helicobasidium*: a unique mycoparasitic-phytoparasitic life strategy. *Mycological Research* **108**: 227–238.
- Maier W, Begerow D, Weiss M, Oberwinkler F (2003) Phylogeny of the rust fungi: an approach using nuclear large subunit ribosomal DNA sequences. *Canadian Journal Botany* **81**: 12–23.
- Marvanová L, Suberkropp K (1990) *Camptobasidium hydrophilum* and its anamorph, *Crucella subtilis*: a new heterobasidiomycete from streams. *Mycologia* **82**: 208–217.
- Mathre DE (1996) Dwarf bunt: politics, identification, and biology. *Annual Review of Phytopathology* **34**: 67–86.
- McNeill J, Barrie FF, Buck WR, Demoulin V, Greuter W, et al. (eds) (2012) *International Code of Nomenclature for algae, fungi, and plants (Melbourne Code)*. [Regnum vegetabile no. 154.] Königstein: Koeltz Scientific Books.
- McTaggart AR, Shivas RG, Boekhout T, Oberwinkler F, Vánky K et al. (2016). *Mycosarcoma (Ustilaginaceae)*, a resurrected generic name for corn smut (*Ustilago maydis*) and its close relatives with hypertrophied, tubular sori. *Journal of Plant Pathology* **7**: 309–315.
- Meinecke EP (1920) Facultative heteroecism in *Peridermium cerebrum* and *Peridermium harknessii*. *Phytopathology* **10**: 279–297.
- Mordue JEM (1988) *Thecaphora solani*. *Descriptions of Pathogenic Fungi and Bacteria* **966**: 1–2.
- Mordue JEM, Gibson IAS (1978) *Cronartium flaccidum*. *Descriptions of Pathogenic Fungi and Bacteria* **580**: 1–2.
- Oberwinkler F, Bandoni RJ (1982) A taxonomic survey of the gasteroid, auricularioid *Heterobasidiomycetes*. *Canadian Journal of Botany* **60**: 1726–1750.
- Oberwinkler F, Bauer R, Bandoni RJ (1990) *Heterogastridiales*: a new order of *Basidiomycetes*. *Mycologia* **82**: 48–58.
- Oberwinkler F, Kirschner R, Arenal F, Villarreal M, Ruio V, et al. (2006) Two new pycnidial members of the *Atractiellales*: *Basidiopycnis hyalina* and *Proceropycnis pinicola*. *Mycologia* **98**: 637–649.
- Ono Y (2015) *Kuehneola* species (*Phragmidiaceae*, *Pucciniales*) on *Vitaceae* plants. *Mycological Progress* **14**: 50.
- Ono Y, Kakishima M, Kudo A, Sato S (1986) *Blastospora smilacis*, a teleomorph of *Caeoma makinoi*, and its sorus development. *Mycologia* **78**: 253–262.
- Ono Y, Kakishima M, Sato S, Harada Y (1987) *Blastospora itoana* on *Smilax nipponica* forms a *Caeoma* stage on *Prunus grayana* in Japan. *Mycologia* **79**: 668–673.
- Petersen RH (1975) Report of the Special Committee for Fungi and Lichens. *Taxon* **24**: 533–534.
- Quandt CA, Kepler RM, Gams W, Araújo JPM, Ban S, et al. (2014) Phylogenetic-based nomenclatural proposals for *Ophiocordycipitaceae* (Hypocreales) with new combinations in *Tolypocladium*. *IMA Fungus* **5**: 121–134.
- Réblová M, Miller AN, Rossman AY, Seifert KA, Crous PW (2016) Recommendations for competing sexual-asexually typified generic names in *Sordariomycetes* (except *Diaporthales*, *Hypocreales*, and *Magnaporthales*). *IMA Fungus* **7**: 131–153.
- Ritz CM, Maier WF, Oberwinkler F, Wissemann V (2005) Different evolutionary histories of two *Phragmidium* species infecting the same dog rose hosts. *Mycological Research* **109**: 603–609.
- Roberts P (1999) *Rhizoctonia-forming fungi: a taxonomic guide*. Kew: Royal Botanic Gardens.
- Rogers DP (1953) Disposition of nomina generica conservanda proposita for fungi. *Mycologia* **45**: 312–322.
- Rossman AY, Adams GC, Cannon PF, Castlebury LA, Crous PW, et al. (2015a) Recommendations of generic names in *Diaporthales* competing for protection or use. *IMA Fungus* **6**: 145–154.
- Rossman AY, Allen WC, Braun U, Castlebury LA, Crous PW, et al. (2016) Overlooked competing asexual and sexually typified generic names of *Ascomycota* with recommendations for their use or protection. *IMA Fungus* **7**: 289–308.

- Rossmann AY, Crous PW, Hyde KD, Hawksworth DL, Aptroot A, *et al.* (2015b) Recommended names of pleomorphic genera in *Dothideomycetes*. *IMA Fungus* **6**: 507–523.
- Rossmann AY, Seifert KA, Samuels GJ, Minnis AM, Schroers H-J, *et al.* (2013) Genera in *Bionectriaceae*, *Hypocreaceae*, and *Nectriaceae* (*Hypocreales*) proposed for acceptance or rejection. *IMA Fungus* **4**: 41–51.
- Salazar-Yepes M, Carvalho jr AA, Hennen JF (2012) *Blastospora* in the Americas: a new combination and anamorphic genus and species. *Mycologia* **104**: 115–120.
- Salazar-Yepes M, Pardo Cardona VM, Buriticá Céspedes P (2007) Especies de Colombia, Ecuador y Perú pertenecientes al género *Gerwasia* Raciborski del Orden Uredinales. *Caldasia* **29**: 105–120.
- Sampaio JP (2011) *Sporidiobolus* Nyland (1949). In: *The Yeasts: a taxonomic study* (Kurtzman CP, Fell JW, Boekhout T, eds): 1549–1561. 5th edn. Amsterdam: Elsevier.
- Samson RA, Visagie CM, Houbraken J, Hong S-B, Hubka V, *et al.* (2014) Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology* **78**: 141–173.
- Sayama A, Kobayashi K, Ogoshi A (1994) Morphological and physiological comparisons of *Helicobasidium mompa* and *H. purpureum*. *Mycoscience* **35**: 15–20.
- Scholz H, Scholz I (1988) Taxonomische Erkenntnisse bei der Erarbeitung einer Brandpilzflora von Deutschland (*Ustilaginales*). *Nova Hedwigia* **47**: 69–79.
- Schumann G (1991) *Plant Diseases: their biology and social impact*. St Paul, MN: American Phytopathological Society Press.
- Simpson JA, Thomas K, Grgurinovic CA (2006) *Uredinales* species pathogenic on species of *Myrtaceae*. *Australasian Plant Pathology* **35**: 549–562.
- Smith PA (2015) Rust fungi on ferns. *Field Mycology* **16**: 49–53.
- Stadler M, Kuhnert E, Peršoh D, Fournier J (2013) The *Xylariaceae* as model example for a unified nomenclature following the “One fungus-one name” (1F1N) concept. *Mycology* **4**: 5–21.
- Statzell-Tallman A, Fell J (1998) 105-*Rhodotorula*. In: *The Yeasts: a taxonomic study* (Kurtzman CP, Fell JW, eds): 800–827. 4th edn. Amsterdam: Elsevier.
- Tanase C, Negrean G (2007) The genus *Milesina* Magnus (*Fungi*, *Basidiomycota*) in Romania. *Analele Stiintifice ale Universitatii “Al. I. Cuza” din Iasi, Biologie Vegetala, ser. nov., Sect. II A* **53**: 108–113.
- Thines M (2016) (2467) Proposal to conserve the name *Ustilago* (*Basidiomycota*) with a conserved type. *Taxon* **65**: 1170–1171.
- Thirumalachar MJ, Kern FD, Patil BV (1966) *Elateraecium*, a new form genus of *Uredinales*. *Mycologia* **58**: 391–396.
- Thirumalachar MJ, Kern FD, Patil BV (1975) *Hiratsukamyces*, a new genus of the *Pucciniastreae*. *Sydowia* **27**: 78–81.
- Toome M, Aime MC (2014) *Pycnopulvinus aurantiacus*, gen. & sp. nov., a new sporocarp-forming member of the *Pucciniomycotina* from Ecuador. *MycoKeys* **8**: 43–50.
- Toome M, Pennycook SR, Aime MC (2015) The generic names *Kriegeria*. *Mycotaxon* **130**: 321–328.
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, *et al.* (eds) (2018) *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017*. [Regnum Vegetabile no. 159.] Königstein: Koeltz Botanical Books.
- Turland NJ, Wiersema JH, Munro AM, Deng Y-F, Zhang L (2017) XIX International Botanical Congress: report of Congress action on nomenclature proposals. *Taxon* **66**: 1234–1245.
- Van der Walt JP (1970) The perfect and imperfect states of *Sporobolomyces salmonicolor*. *Antonie van Leeuwenhoek* **36**: 49–55.
- Van der Walt JP (1973) *Aessosporon dendrophilum* sp. nov., the perfect state of *Bullera dendrophila*. *Antonie van Leeuwenhoek* **39**: 455–460.
- Vánky K (2012) *Smut Fungi of the World*. St Paul, MN: American Phytopathological Society Press.
- Vánky, K, Lutz M, Bauer R (2008) About the genus *Thecaphora* (*Glomosporiaceae*) and its new synonyms. *Mycological Progress* **7**: 31–39.
- Visagie CM, Houbraken J, Frisvad JC, Hong S-B, Klaassen CHW, *et al.* (2014) Identification and nomenclature of the genus *Penicillium*. *Studies in Mycology* **78**: 343–371.
- Vogler DR, Bruns TD (1998) Phylogenetic relationships among the pine stem rust fungi (*Cronartium* and *Peridermium* spp.). *Mycologia* **90**: 244–257.
- Wakefield E (1939) Nomina generica conservanda: Contributions from the nomenclature committee of the British Mycological Society. I. *Transactions of the British Mycological Society* **23**: 215–232.
- Wang Q-M, Yurkov AM, Göker M, Lumbsch HT, Leavitt SD, *et al.* (2015a) Phylogenetic classification of yeasts and related taxa within *Pucciniomycotina*. *Studies in Mycology* **81**: 149–189.
- Wang Q-M, Begerow D, Groenewald M *et al.* (2015b) Multigene phylogeny and taxonomic revision of yeasts and related fungi in the *Ustilaginomycotina*. *Studies in Mycology* **81**: 27–53.
- Wijayawardene DNN, McKenzie EHC, Hyde KD (2012) Towards incorporating anamorphic fungi in a natural classification – checklist and notes for 2011. *Mycosphere* **3**: 157–228.
- Wingfield BD, Ericson L, Szaro T, Burdon JJ (2004) Phylogenetic patterns in the *Uredinales*. *Australasian Plant Pathology* **33**: 327–335.
- Wright JE (1970) *Agaricostilbum*, a new genus of *Deuteromycetes* on palm spathes from Argentina. *Mycologia* **62**: 679–682.
- Wright JE, Bandoni R, Oberwinkler F (1981) *Agaricostilbum*: an auricularioid basidiomycete. *Mycologia* **73**: 880–886.
- Yun HY, Hong SG, Rossmann AY, Lee SK, Lee KJ, *et al.* (2009) The rust fungus *Gymnosporangium* in Korea including two new species, *G. monticola* and *G. unicolorne*. *Mycologia* **101**: 790–809.
- Yun H-Y, Minnis AM, Kim Y-H, Castlebury LA, Aime MC (2011) The rust genus *Frommeëlla* revisited: a later synonym of *Phragmidium* after all. *Mycologia* **103**: 1451–1463.
- Zhang N, Luo J, Rossmann A, Aoki T, Chuma I, *et al.* (2016) Generic names in *Magnaporthales*. *IMA Fungus* **7**: 155–159.